

## Does a physiological role for KCNE subunits exist in the immune system?

Laura Solé & Antonio Felipe

To cite this article: Laura Solé & Antonio Felipe (2010) Does a physiological role for KCNE subunits exist in the immune system?, Communicative & Integrative Biology, 3:2, 166-168, DOI: [10.4161/cib.3.2.10602](https://doi.org/10.4161/cib.3.2.10602)

To link to this article: <https://doi.org/10.4161/cib.3.2.10602>



© 2010 Landes Bioscience



Published online: 01 Mar 2010.



Submit your article to this journal [↗](#)



Article views: 192



View related articles [↗](#)



Citing articles: 6 View citing articles [↗](#)

## Does a physiological role for KCNE subunits exist in the immune system?

Laura Solé and Antonio Felipe\*

Molecular Physiology Laboratory; Departament de Bioquímica i Biologia Molecular; Institut de Biomedicina; Universitat de Barcelona; Barcelona, Spain

The study of channel modulation by regulatory subunits has attracted considerable attention. Evidence indicates a pivotal role for accessory proteins in the channelosome. For instance, these regulatory subunits are necessary to recapitulate *in vivo* ion currents and to further understand the physiological role of ion channels. KCNEs are a family of regulatory subunits that interact with a wide range of channels. We have described for the first time a molecular interaction between KCNE4 and the voltage-dependent potassium channel  $K_v1.3$ . The association of KCNE4, which alters the biophysical properties, trafficking and membrane localization of  $K_v1.3$ , functions as an endogenous dominant-negative mechanism. Since both proteins are expressed in the immune system,  $K_v1.3$ /KCNE4 channels may contribute to the fine-tuning of the immune response. Therefore, our results point to KCNE4 as a novel target for immunomodulation. KCNE4 is not the only KCNE which is expressed in leukocytes. All KCNEs (KCNE1-5) are present, and some members demonstrate modulation during proliferation and cancer. In summary, regulatory KCNE subunits are expressed in the immune system. In addition, several voltage-dependent  $K^+$  channels, which could interact with KCNEs, are also detected. Therefore, KCNE subunits may play a yet undiscovered role in the physiology of the immune system.

modulate a large number of voltage-dependent  $K^+$  channels.<sup>1</sup> The most well characterized interaction occurs with KCNQ1 ( $K_v7.1$ ) channels.<sup>2-4</sup>  $K_v7.1$ /KCNE1 channels recapitulate the cardiac  $I_{ks}$  current.<sup>5</sup> Several KCNE1 mutations, which trigger severe cardiac channelopathies, demonstrate the pivotal function of KCNE1 on cardiovascular physiology.<sup>6-8</sup> The implication of that the remaining KCNE peptides may contribute to the modulation of  $I_{ks}$  is now under intense investigation.<sup>9-13</sup> Besides  $K_v7.1$ , KCNE members associate with other  $K^+$  channels. Thus,  $K_v11.1$  in association with KCNE2 conducts the cardiac  $I_{kr}$  current.<sup>14</sup>

Although many KCNE subunits share tissue expression with Shaker  $K^+$  channels ( $K_v1$ ) channels, KCNE interactions with  $K_v1$  channels have attracted little research.<sup>1</sup> Early in 1992,  $K_v1.3$  and KCNE1 were simultaneously cloned in human Jurkat T-cells.<sup>15</sup> Ten years later, Grunnet et al. (2003) described that KCNE4 modulates  $K_v1.1$  and  $K_v1.3$  channels and their heteromeric forms. However, while  $K_v1.1$  and  $K_v1.3$  biophysics were affected,  $K_v1.1$  membrane surface targeting was not altered.<sup>16</sup> Later, Melman et al. (2004) described that KCNE1 co-immunoprecipitated with  $K_v1.5$ , but no further research was undertaken.<sup>17</sup> Recently, Abbot et al. (2008) demonstrated that *kcne2*<sup>-/-</sup> mice exhibit altered cardiac  $I_{kslow}$ , identified an interaction between  $K_v1.5$  and KCNE2 in murine ventricles and suggested a functional role for KCNE2 in promoting  $K_v1.5$  surface expression.<sup>18</sup>

In this context, our recent contribution unequivocally demonstrates molecular interactions between KCNE4 and  $K_v1.3$ ,

**Key words:** voltage-dependent potassium channels, KCNE4,  $K_v1.3$ , leukocytes, regulatory subunits

Submitted: 11/10/09


Accepted: 11/10/09

Previously published online:  
[www.landesbioscience.com/journals/cib/article/10602](http://www.landesbioscience.com/journals/cib/article/10602)


\*Correspondence to: Antonio Felipe;  
Email: [afelipe@ub.edu](mailto:afelipe@ub.edu)

Addendum to: Solé L, Roura-Ferrer M, Pérez-Verdaguer M, Oliveras A, Calvo M, Fernández-Fernández JM, Felipe A. KCNE4 suppresses  $K_v1.3$  currents by modulating trafficking, surface expression and channel gating. *J Cell Sci* 2009; 122:3738–48; PMID: 19773357; DOI:10.1242/jcs.056689.

KCNEs are a group of regulatory subunits composed of 5 members (KCNE1-5). KCNE peptides are small single spanning membrane proteins (<20 kDa) which



**Figure 1.** Human KCNE expression in healthy and cancer tissues. Affymetrix GeneChips HG-U95A-E (GeneNote, [http://bioinfo2.weizmann.ac.il/cgi-bin/genenote/home\\_page.pl](http://bioinfo2.weizmann.ac.il/cgi-bin/genenote/home_page.pl)) and HG-UI33A (GNF, (<http://biogps.gnf.org>) normalized as described in GeneCard (<http://www.genecards.org>). Black columns: healthy tissue (GeneNote); Grey columns: healthy tissues (GNF); White columns: cancer samples (GNF).



**Figure 2.** KCNE expression in human leukocytic cell lines. Array data from GNF BioGPS (<http://biogps.gnf.org>) normalized according to Su et al.<sup>31</sup> Legend: CD34, bone marrow CD34<sup>+</sup> progenitors; CD105, endothelial; X212, B lymphoblasts; CD19, B cells; BDCA4, dendritic cells; CD8, CD8<sup>+</sup> T-cells; CD4, CD4<sup>+</sup> T-cells; CD56, NK cells; CD33, myeloid; CD14, monocytes.

which impairs the trafficking and localization of these channels.<sup>19</sup> KCNE4 acts as an endogenous dominant-negative regulatory subunit. Further inhibitory mechanisms by KCNE4 have been observed. Grunnet et al. (2002) describe that KCNE4 abrogates  $K_{V7.1}$  currents.<sup>20</sup> Later, George and coworkers (2008) postulate that KCNE4 could form part of a  $K_{V7.1}$ /KCNE1 heterocomplex, responsible for downregulating the  $I_{Ks}$  current.<sup>10</sup> Furthermore, KCNE4 also inhibits the calcium-activated K<sup>+</sup> channel ( $K_{Ca1.1}$ )<sup>21</sup> and oligomeric  $K_{V4.2}$  + KChIP2 channels.<sup>22</sup>

Although limited, the immunitary system has a defined repertoire of K<sup>+</sup> channel genes. Leukocytes express,  $K_{V1.3}$ ,  $K_{V1.5}$  and several  $K_{V\beta}$  regulatory subunits.<sup>23-25</sup>

We have demonstrated that  $K_{V1.3}$  and  $K_{V1.5}$  form heteromeric complexes to fine-tune the immunitary response.<sup>23,26</sup> The evidence that KCNE4 is also implicated,<sup>19</sup> the presence of KCNE1 in lymphocytes,<sup>15</sup> and its putative association to  $K_{V1.5}$ ,<sup>17</sup> suggest an unidentified role for KCNE peptides in immunitary system physiology.

In fact, our recent study is not the first to describe the presence of the KCNEs in the immunitary system. KCNE1 was cloned in T-lymphocytes.<sup>15</sup> Grunnet et al. (2003) found KCNE4 mRNA in lymphocytes.<sup>16</sup> In addition, KCNEs mRNA expression has also been investigated in thymus.<sup>10,27</sup> Here we recapitulate the information available about the expression of KCNE1-5 in different tissue arrays, all

derived from the immune system. To that end, data from tissues and tumors, cell lines and cancer cell lines are shown in **Figures 1, 2 and 3**, respectively. All KCNE1-5 members have been detected in myeloid and lymphoid lineages. It is noteworthy that while KCNE1 expression increases in some cancers, KCNE2 and KCNE4 decrease. Similarly, KCNE1 induction has also been detected in germinal tumors.<sup>28</sup> We have previously described the importance of  $K_{V1.3}$  during activation and proliferation of macrophages.<sup>25,26</sup> Although both processes augment  $K_{V1.3}$ , the channel seems to play a dual role. Prolonged signaling during activation triggers apoptosis.<sup>29,30</sup> Additionally, highly proliferative cells exhibit lower KCNE4 expression

than their counterparts do (Fig. 1). During an insult,  $K_v1.3$  is activated and sustained activation triggers cell death. The persistence of activated macrophages during inflammation leads to disease. In this scenario, KCNE4 would exert an inhibitory effect not needed in proliferation. Accordingly, it is worth noting that during proliferation, most cells become resistant to apoptosis.


Evidence demonstrates that KCNE subunits are present in the immune system and may interact with  $K_v1$  channels. Therefore, we suggest that KCNE subunits may play a yet undiscovered role in the immune system via associations with leukocyte  $K^+$  channels. These new interactions situate KCNE, specifically KCNE4, as novel targets for immunomodulation. Further research should be undertaken to shed some light on this new issue.

### Acknowledgements

The authors thank to all present and past members of the molecular physiology laboratory. Supported by the Ministerio de Ciencia e Innovación (MICINN), Spain (BFU2008-00431 and CSD2008-00005). L.S. is a fellow from MICINN.

### References

1. McCrossan ZA, Abbott GW. The MinK-related peptides. *Neuropharmacology* 2004; 47:787-821.
2. Haitin Y, Wiener R, Shaham D, Peretz A, Cohen EB, Shamgar L, et al. Intracellular domains interactions and gated motions of I(KS) potassium channel subunits. *EMBO J* 2009; 28:1994-2005.
3. Chen J, Zheng R, Melman YF, McDonald TV. Functional interactions between KCNE1 C-terminus and the KCNQ1 channel. *PLoS One* 2009; 4:5143.
4. Kang C, Tian C, Sonnichsen FD, Smith JA, Meiler J, George AL Jr, et al. Structure of KCNE1 and implications for how it modulates the KCNQ1 potassium channel. *Biochemistry* 2008; 47:7999-8006.
5. Barhanin J, Lesage F, Guillemare E, Fink M, Lazdunski M, Romey G. K(V)LQT1 and IsK (minK) proteins associate to form the I(Ks) cardiac potassium current. *Nature* 1996; 384:78-80.
6. Schulze-Bahr E, Wang Q, Wedekind H, Haverkamp W, Chen Q, Sun Y, et al. KCNE1 mutations cause jervell and Lange-Nielsen syndrome. *Nat Genet* 1997; 17:267-8.
7. Splawski I, Tristani-Firouzi M, Lehmann MH, Sanguinetti MC, Keating MT. Mutations in the hminK gene cause long QT syndrome and suppress I(Ks) function. *Nat Genet* 1997; 17:338-40.
8. Chiang CE, Roden DM. The long QT syndromes: genetic basis and clinical implications. *J Am Coll Cardiol* 2000; 36:1-12.
9. Lundquist AL, Manderfield LJ, Vanoye CG, Rogers CS, Donahue BS, Chang PA, et al. Expression of multiple KCNE genes in human heart may enable variable modulation of I(Ks). *J Mol Cell Cardiol* 2005; 38:277-87.



**Figure 3.** KCNE expression in different human leukemia and lymphoma cell lines. Array data from GNF BioGPS (<http://biogps.gnf.org>) normalized according to Su et al.<sup>31</sup> Legend: MOLT4, Lymphoblastic Leukemia; K.582, Chronic Myelogenous Leukemia; Daudi, Burkitt's Lymphoma; HL60, Promyelocytic Leukemia; Raji, Burkitt's Lymphoma.

10. Manderfield LJ, George AL Jr. KCNE4 can co-associate with the I(Ks) (KCNQ1-KCNE1) channel complex. *Febs J* 2008; 275:1336-49.
11. Tseng GN. The phenotype of a KCNQ1 mutation depends on its KCNE partners: is the cardiac slow delayed rectifier (IKs) channel more than a KCNQ1/KCNE1 complex? *Heart Rhythm* 2007; 4:1542-3.
12. Morin TJ, Kobertz WR. A derivatized scorpion toxin reveals the functional output of heteromeric KCNQ1-KCNE K<sup>+</sup> channel complexes. *ACS Chem Biol* 2007; 2:469-73.
13. Bendahhou S, Marionneau C, Haurogue K, Larroque MM, Derand R, Szuts V, et al. In vitro molecular interactions and distribution of KCNE family with KCNQ1 in the human heart. *Cardiovasc Res* 2005; 67:529-38.
14. Abbott GW, Sesti F, Splawski I, Buck ME, Lehmann MH, Timothy KW, et al. MiRP1 forms IKr potassium channels with HERG and is associated with cardiac arrhythmia. *Cell* 1999; 97:175-87.
15. Attali B, Romey G, Honore E, Schmid-Alliana A, Mattei MG, Lesage F, et al. Cloning, functional expression, and regulation of two K<sup>+</sup> channels in human T lymphocytes. *J Biol Chem* 1992; 267:8650-7.
16. Grunnet M, Rasmussen HB, Hay-Schmidt A, Rosenstierne M, Klaerke DA, Olesen SP, Jespersen T. KCNE4 is an inhibitory subunit to K<sub>v</sub>1.1 and K<sub>v</sub>1.3 potassium channels. *Biophys J* 2003; 85:1525-37.
17. Melman YF, Um SY, Krummerman A, Kagan A, McDonald TV. KCNE1 binds to the KCNQ1 pore to regulate potassium channel activity. *Neuron* 2004; 42:927-37.
18. Roepke TK, Kontogeorgis A, Ovanec C, Xu X, Young JB, Purtell K, et al. Targeted deletion of *kcne2* impairs ventricular repolarization via disruption of I(Ks,slow) and I(to,f). *FASEB J* 2008; 22:3648-60.
19. Sole L, Roura-Ferrer M, Perez-Verdaguer M, Oliveras A, Calvo M, Fernandez-Fernandez JM, Felipe A. KCNE4 suppresses K<sub>v</sub>1.3 currents by modulating trafficking, surface expression and channel gating. *J Cell Sci* 2009; 122:3738-48.
20. Grunnet M, Jespersen T, Rasmussen HB, Ljungstrom T, Jorgensen NK, Olesen SP, Klaerke DA. KCNE4 is an inhibitory subunit to the KCNQ1 channel. *J Physiol* 2002; 542:119-30.
21. Levy DI, Wanderling S, Biemesderfer D, Goldstein SA. MiRP3 acts as an accessory subunit with the BK potassium channel. *Am J Physiol Renal Physiol* 2008; 295:380-7.
22. Radicke S, Cotella D, Graf EM, Banse U, Jost N, Varro A, et al. Functional modulation of the transient outward current I<sub>to</sub> by KCNE beta-subunits and regional distribution in human non-failing and failing hearts. *Cardiovasc Res* 2006; 71:695-703.
23. Vicente R, Escalada A, Villalonga N, Texido L, Roura-Ferrer M, Martin-Satue M, et al. Association of K<sub>v</sub>1.5 and K<sub>v</sub>1.3 contributes to the major voltage-dependent K<sup>+</sup> channel in macrophages. *J Biol Chem* 2006; 281:37675-85.
24. Vicente R, Escalada A, Soler C, Grande M, Celada A, Tamkun MM, et al. Pattern of K<sub>v</sub> beta subunit expression in macrophages depends upon proliferation and the mode of activation. *J Immunol* 2005; 174:4736-44.
25. Vicente R, Escalada A, Coma M, Fuster G, Sanchez-Tillo E, Lopez-Iglesias C, et al. Differential voltage-dependent K<sup>+</sup> channel responses during proliferation and activation in macrophages. *J Biol Chem* 2003; 278:46307-20.
26. Villalonga N, Escalada A, Vicente R, Sanchez-Tillo E, Celada A, Solsona C, Felipe A. K<sub>v</sub>1.3/K<sub>v</sub>1.5 heteromeric channels compromise pharmacological responses in macrophages. *Biochem Biophys Res Commun* 2007; 352:913-8.
27. Lundquist AL, Turner CL, Ballester LY, George AL Jr. Expression and transcriptional control of human KCNE genes. *Genomics* 2006; 87:119-28.
28. Tsevi I, Vicente R, Grande M, Lopez-Iglesias C, Figueras A, Capella G, et al. KCNQ1/KCNE1 channels during germ-cell differentiation in the rat: expression associated with testis pathologies. *J Cell Physiol* 2005; 202:400-10.
29. Detre C, Kiss E, Varga Z, Ludanyi K, Paszty K, Enyedi A, et al. Death or survival: membrane ceramide controls the fate and activation of antigen-specific T-cells depending on signal strength and duration. *Cell Signal* 2006; 18:294-306.
30. Szabo I, Adams C, Gullbins E. Ion channels and membrane rafts in apoptosis. *Pflügers Arch* 2004; 448:304-12.
31. Su AI, Wiltshire T, Batalov S, Lapp H, Ching KA, Block D, et al. A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc Natl Acad Sci USA* 2004; 101:6062-7.